

Linear algebra meets Lie algebra: the Kostant–Wallach theory

Noam Shomron
Beresford N. Parlett*

22nd October 2008

Abstract

In two languages, Linear Algebra and Lie Algebra, we describe the results of Kostant and Wallach on the fibre of matrices with prescribed eigenvalues of all leading principal submatrices. In addition, we present a brief introduction to basic notions in Algebraic Geometry, Integrable Systems, and Lie Algebra aimed at specialists in Linear Algebra.

0 Introduction

The well-known meeting-place of linear algebra and Lie algebra is the classical matrix groups. This paper is not a survey of that theory, but is about a more specific confluence, and we consider only the general linear (“type a”) case among the classical groups.

For want of a standard name we designate as the Ritz values of a matrix the eigenvalues of all its leading principal submatrices. In [KW1] and [KW2], Kostant and Wallach studied the structure of the set (“fibre”) of matrices with given Ritz values. They constructed a certain commutative Lie group, which acts on the space of matrices and whose action preserves Ritz values; the second paper culminates by showing that this naturally leads to a particularly nice set of coordinates on the space of those matrices whose Ritz values satisfy some disjointness condition.

Inspired by this work, Parlett and Strang [PS] studied such problems using bona fide matrix theory and linear algebra. Later one of us [BNP] showed quite explicitly how to parameterize the space of matrices with given generic Ritz values, without invoking any Lie theory or algebraic geometry. However, hiding away the symmetry of the problem does have some drawbacks: while the coordinates are easy to define, it is not clear what they mean, or that they satisfy any natural properties. Thus the extra structure of Lie theory can give depth to the matrix theory.

*Mathematics Department and Computer Science Division in the EECS Department, University of California, Berkeley, Calif., 94720 USA. Email: parlett@math.berkeley.edu

Kostant and Wallach's group, and their parameterization of generic fibres, do appear as such in the matrix-theoretic approach, but the properties (such as their version of the Gelfand–Kirillov theorem) that mark them as nice can only be seen by considering the global geometry of the space of matrices, rather than a single fibre at a time.

The purpose of the present work is twofold. The primary purpose is expository. We need to introduce enough of the language of Lie theory to be able to state and apply some of Kostant and Wallach's results. We hope the reader will be convinced that geometrical intuition, using the machinery of Lie/algebraic group theory and algebraic geometry, while on the surface very abstract, can not only suggest the right way to think about a problem in linear algebra, but, in fact, tell us how to do the actual computation.

We show how to recover BNP's construction using the language and results of Kostant and Wallach, and prove that the two sets of coordinates are, in fact, identical.

1 Matrix picture

1.1 Notation and basic facts

In contrast to most papers on matrix theory, certain matrices will be denoted by lower-case Roman letters, such as x , for compatibility with the notation used in [KW1, KW2]. However, sometimes lower-case Roman letters, such as b and c , denote vectors, and sometimes we will use them to denote scalars, such as t , etc.; the type of object will always be unambiguous. For a square matrix x , the leading principal submatrix of order m will be denoted by x_m ; in Matlab notation $x_m = x(1:m, 1:m)$.

Let $E(x)$ denote the multiset of eigenvalues of x . The object of study is $\mathbb{C}^{n \times n}$ for a fixed natural number n , but, since it will be endowed with extra (e.g., Lie) structure, we use the standard notation $M(n)$. What is not standard is $\mathcal{R}(x)$, $x \in M(n)$.

Definition. The set of Ritz values of $x \in M(n)$ is the tuple $\mathcal{R}(x) = (E(x_1), E(x_2), \dots, E(x_n))$.

This name was chosen because, in numerical linear algebra, when x is Hermitian $E(x_m)$, for $m < n$, is regarded as an approximation to a subset of $E(x_n)$, and, independently, Rayleigh and Ritz showed that the former are optimal approximations (in various senses) from the subspace spanned by the first m columns of the identity matrix. See [P, Chapter 11].

For Hermitian matrices there are interlacing conditions connecting $E(x_{m-1})$ and $E(x_m)$. However, for $M(n)$, there are no constraints on $\mathcal{R}(x)$; any set of $\binom{n+1}{2}$ complex numbers is $\mathcal{R}(y)$ for some $y \in M(n)$. Moreover, sharing Ritz values determines an equivalence relation on $M(n)$, and we have

$$M(n) = \coprod_{\mathcal{R}} M_{\mathcal{R}}(n), \quad (1.1.1)$$

where $M_{\mathcal{R}}(n) = \{x \in M(n) \mid \mathcal{R}(x) = \mathcal{R}\}$. (The coproduct symbol $\coprod$ here just means the set-theoretic disjoint union.) In geometric theory the equivalence class $M_{\mathcal{R}}(n)$ is called a fibre,¹ and we will use this terminology. We have the following

¹Specifically, a fibre of the map $x \mapsto \mathcal{R}(x)$ which assigns to each matrix its Ritz values.

Problem 1.1.2. Given $\mathcal{R}$, describe the fibre $M_{\mathcal{R}}(n)$.

The first observation is that $\mathcal{R}$ determines the diagonal entries uniquely:

$$x_{mm} = \sum E(x_m) - \sum E(x_{m-1}).$$

Thus all members of $M_{\mathcal{R}}(n)$ share the same diagonal.

Elementary conjugations. For matrix theorists it seems a challenge to generate the fibre for a given $\mathcal{R}$. In the generic case all elements are similar to each other, and yet the diagonal is fixed. What mappings $x \mapsto gxg^{-1}$, with $g \in \mathrm{GL}(n)$, preserve $\mathcal{R}$?

A little reflection suggests two types which we will call elementary conjugations:

- (i) transposition (not conjugate transpose), $x \mapsto x^T$,
- (ii) diagonal similarity, $x \mapsto dxd^{-1}$, $d \in \mathrm{GL}(n)$ diagonal.

These two are far too weak to generate the fibre; the effect of elementary conjugations upon the dual coordinates s which parameterize the fibre will be shown later.

To state the first significant result of Kostant and Wallach, we need the notion of a Hessenberg matrix.

Definition. A matrix $H \in M(n)$ is upper Hessenberg if $H_{ij} = 0$ for $i > j + 1$. H is said to be unreduced if $H_{i+1,i} \neq 0$, $1 \leq i \leq n-1$. H is said to be unit upper Hessenberg if $H_{i+1,i} = 1$, $1 \leq i \leq n-1$.

The result of Kostant and Wallach is that upper Hessenberg matrices serve as a natural set of representatives of each $M_{\mathcal{R}}(n)$. Formally,

Theorem 1.1.3 ([KW1], Theorem 0.1 and Remark 0.3). For any $\mathcal{R}$, $M_{\mathcal{R}}(n)$ contains exactly one unit upper Hessenberg matrix.

Remark. A matrix-oriented proof was given in [PS].

Another result of Kostant and Wallach, also established by elementary means in [PS], is that when $\mathcal{R}$ is generic (defined below in Definition 1.2.1) then the strictly lower triangular part of $x \in M_{\mathcal{R}}(n)$ determines uniquely the strictly upper triangular part, and vice versa. Thus it is tempting to think of the strictly lower part as a suitable set of coordinates for x that is dual, or complementary, to $\mathcal{R}$. The parameter count $\binom{n}{2}$ is exactly right. For reasons that will be made clear below, this temptation must be resisted.

The major result of [KW2] was to find a “nice” set of coordinates to specify the members of $M_{\mathcal{R}}(n)$ for generic $\mathcal{R}$. They are given by tuples $s = (s^{(1)}, \dots, s^{(n-1)})$, with $s^{(m)} \in \mathbb{C}^m$, but no entry can vanish, so we invoke $\mathbb{C}^\times$, the multiplicative group $\mathbb{C} \setminus \{0\}$, and have $s^{(m)} \in (\mathbb{C}^\times)^m$. Thus Kostant and Wallach present a coordinate system $(\mathcal{R}, s)$ for the generic elements of $M(n)$ that is not familiar to matrix theorists. The goal of this paper is to show the geometric meaning of those coordinates. In some sense this is an instance of the Darboux coordinates (q, p) in the Hamilton–Jacobi theory of mechanics.²

²The coordinates $(\mathcal{R}, s)$ will be action-angle coordinates arising from an integrable system; see Sec-

1.2 Eigenvalue disjointness

The simplest version of the theory, on which we will focus and for reasons we will explain, occurs in the “generic” case. Consider the following conditions on an $n \times n$ matrix x :

(G1 _{m}) The elements of $E(x_m)$ are distinct.

(G2 _{m}) $E(x_m) \cap E(x_{m+1}) = \emptyset$.

The significance of these conditions will be discussed in Section 1.5.

Definition 1.2.1. We call (G1 _{m}) and (G2 _{m}) the “eigenvalue disjointness conditions.” If both (G1 _{m}), $1 \leq m \leq n$, and (G2 _{m}), $1 \leq m \leq n - 1$, hold for $x \in M(n)$, we will call x generic.

Definition.

$$M_\Omega(n) = \{ x \in M(n) \mid x \text{ is generic} \}.$$

The complement of $M_\Omega(n)$ in $M(n)$ breaks up into pieces specified by how badly conditions (G1 _{m}) and (G2 _{m}) are violated. Each such violation translates into the vanishing of some polynomial in the entries of x , e.g., (G2₁) is false exactly when $x_{12}x_{21} = 0$, and (G1₂) is false exactly when $(x_{11} - x_{22})^2 + 4x_{12}x_{21} = 0$. The set $M_\Omega(n)$ is, therefore, a (nonempty, therefore dense) Zariski-open subset of $M(n)$.³ For this reason we sometimes say that a matrix $x \in M_\Omega(n)$ has generic Ritz values. Often the term “generic” refers to any dense open subset. For example, one would say the condition that a matrix be diagonalizable is a generic condition. It is somewhat confusing to refer to “generic” matrices, because there are many dense open subsets. In fact, since $M(n) \cong \mathbb{C}^{n^2}$ is an algebraic variety, in the Zariski topology any nonempty open subset is dense. In this paper, for the sake of brevity, “generic” will refer to the specific eigenvalue disjointness conditions just described.

What is wrong with eigenvalues? A given set $\mathcal{R}$ of Ritz values may be designated in various ways by a matrix theorist. We could write down the eigenvalues $E(x_1), E(x_2), \dots, E(x_n)$ in some specific order for each m . We could write down the set $\{P_1, P_2, \dots, P_n\}$ of monic characteristic polynomials of $x_1, x_2, \dots, x_n$. We could write down the coefficients of each P_m other than the dominant one. The descriptions are equivalent. Life is not so carefree for Lie theorists because there is no natural global meaning to “the i th eigenvalue of x .” For example, let

$$x = x(t) = \begin{pmatrix} 0 & e^{2\pi i t} \\ 1 & 0 \end{pmatrix}.$$

As t goes from 0 to 1, $x(t)$ describes a smooth family of generic matrices. We may diagonalize $x(0) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ via

$$\begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}^{-1} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

tion 3.

³If the appellation “Zariski” is intimidating, do not fret. Our exposition eschews further mention of it.

Extending this to the family, we have

$$\begin{pmatrix} e^{\pi it} & -e^{\pi it} \\ 1 & 1 \end{pmatrix}^{-1} x(t) \begin{pmatrix} e^{\pi it} & -e^{\pi it} \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} e^{\pi it} & 0 \\ 0 & -e^{\pi it} \end{pmatrix} =: \Lambda(t).$$

However, $\Lambda(0) \neq \Lambda(1)$ despite the fact that $x(0) = x(1)$. Hence there is no consistent smooth global way to order the eigenvalues of a matrix. But Lie algebra is committed to smooth maps.

In fact, Kostant and Wallach do give a global definition of “ i th eigenvalue” by means of a “covering” $M_\Omega(n, \mathfrak{e})$ of $M_\Omega(n)$. This extra technicality is not needed for a description of the fibres (it is only introduced in the second part [KW2] of their paper which establishes a “Gelfand–Kirillov theorem” for $M(n)$). We avoid this complication by considering a single fibre $M_{\mathcal{R}}(n)$ with some ordering of each $E(x_m)$ already given.

1.3 The complementary coordinates

Now we describe the complementary coordinates $s = (s_1, \dots, s_{\binom{n}{2}})$ for a generic fibre. Consider a matrix x with generic Ritz values. Write $\mathcal{R}(x) = (E(x_1), \dots, E(x_n))$ with a fixed ordering for each $E(x_m)$. We will denote by

$$\Lambda_m = \text{diag}(\mu_1^{(m)}, \dots, \mu_m^{(m)})$$

the diagonal matrix with the elements of $E(x_m)$ placed along the diagonal. For $1 \leq m \leq n-1$, (G1 _{m}) implies that x_m is similar to Λ_m . Hence there exists a matrix $g_m \in \text{GL}(m)$ such that

$$x_m = g_m \Lambda_m g_m^{-1}, \quad (1.3.1)$$

and it becomes unique if the last row of g_m consists of ones. (Note that the last entry of an eigenvector of x_m must not vanish, since $(\lambda I_m - x_m) \begin{pmatrix} u \\ 0 \end{pmatrix} = 0$ implies $(\lambda I_{m-1} - x_{m-1})u = 0$, but x_{m-1} and x_m are assumed to have no eigenvalues in common.) Then

$$x_{m+1} = \begin{pmatrix} g_m & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \Lambda_m & c_m \\ b_m^T & \delta_{m+1} \end{pmatrix} \begin{pmatrix} g_m^{-1} & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} x_m & g_m c_m \\ b_m^T g_m^{-1} & \delta_{m+1} \end{pmatrix} \quad (1.3.2)$$

(we consistently write rows as transposed columns), and our dual coordinates appear in (1.3.2) as the entries of b_m^T . We call the pair (b_m^T, c_m) the “arrow coordinates” of x_{m+1} .

Claim 1.3.3. b_m^T is identical with Kostant and Wallach’s coordinates $s^{(m)}$.

A proof is given in Section 3.1.

Notation. $\text{diag}(v)$, $v \in \mathbb{C}^m$, denotes the diagonal matrix $\text{diag}(v_1, v_2, \dots, v_m) \in M(m)$.

It is a consequence of the generic conditions (G1 _{m}) and (G2 _{m}) that $\text{diag}(b_m) \text{diag}(c_m)$ is invertible:

Theorem 1.3.4. $\text{diag}(b_m) \text{diag}(c_m) = -P_{m+1}(\Lambda_m)(P'_m(\Lambda_m))^{-1}$.

Proof. By (1.3.2),

$$\begin{aligned}
P_{m+1}(\lambda) &= \det(\lambda I_{m+1} - x_{m+1}) \\
&= \det \begin{pmatrix} \lambda I_m - \Lambda_m & -c_m \\ -b_m^T & \lambda - \delta_{m+1} \end{pmatrix} \\
&= P_m(\lambda) [(\lambda - \delta_{m+1}) - b_m^T (\lambda I_m - \Lambda_m)^{-1} c_m] \\
&= P_m(\lambda)(\lambda - \delta_{m+1}) - b_m^T \text{diag} \left(P_m^{(1)}(\lambda), \dots, P_m^{(m)}(\lambda) \right) c_m,
\end{aligned} \tag{1.3.5}$$

where

$$P_m^{(i)}(\lambda) = \prod_{\mu \in E(x_m) \setminus \{\mu_i^{(m)}\}} (\lambda - \mu), \quad P_m^{(i)}(\mu_i^{(m)}) = P'_m(\mu_i^{(m)}).$$

Evaluating (1.3.5) at $\lambda = \mu_1^{(m)}, \mu_2^{(m)}, \dots, \mu_m^{(m)}$ gives

$$P_{m+1}(\Lambda_m) = 0 - \text{diag}(b_m) P'_m(\Lambda_m) \text{diag}(c_m),$$

and diagonal matrices commute. $\square$

Given all b_m^T , we can reconstruct a unique $x \in M_{\mathcal{A}}(n)$. First, a useful lemma:

Lemma 1.3.6. Consider a (down) arrow matrix

$$A = \begin{pmatrix} D & p \\ q^T & \delta \end{pmatrix} \in M(m+1),$$

where $D = \text{diag}(d_i)$ and A is similar to $\Lambda = \text{diag}(\lambda_j)$, and d_i and λ_j are all distinct. It is convenient to define the (rectangular and skew) matrix Cauchy(D, Λ),

$$\text{Cauchy}(D, \Lambda)_{ij} = (d_i - \lambda_j)^{-1}$$

(usually Cauchy matrices are defined as $(d_i + \lambda_j)^{-1}$ with same-sized parameter sets). Also, define **ones** to be an array all of whose entries are 1's; the shape of **ones** is dictated by the context. Then the spectral factorization of A is given by

$$A = Z^{-1} \Lambda Z,$$

where

$$\begin{aligned}
Z^{-1} &= \begin{bmatrix} -\text{diag}(p) \text{Cauchy}(D, \Lambda) \\ \mathbf{ones} \end{bmatrix}, \quad Z = \Pi^{-1} [\text{Cauchy}(\Lambda, D) \text{diag}(q), \mathbf{ones}], \\
\Pi &= -\text{Cauchy}(\Lambda, D) \text{diag}(q) \text{diag}(p) \text{Cauchy}(D, \Lambda) + \mathbf{ones}, \\
\Pi_{ij} &= 0 \text{ if } i \neq j, \quad \Pi_{jj} = 1 + \sum_i \frac{p_i q_i}{(\lambda_j - d_i)^2} = 1 - \sum_i \frac{\prod_m (d_i - \lambda_m)}{(\lambda_j - d_i)^2 \prod_{k \neq i} (d_i - d_k)}, \\
&\quad (\text{thus } \Pi \text{ is independent of } p \text{ and } q)
\end{aligned}$$

i.e., we have found the eigenvectors of A .

Proof. The distinctness of the d_i and λ_j implies that the last entry of an eigenvalue of A must be non-zero. If μ is an eigenvalue of A , the equations

$$(\mu I - A) \begin{pmatrix} u \\ 1 \end{pmatrix} = 0, \quad (v^T \quad 1) (\mu I - A) = 0$$

imply

$$\begin{aligned} u &= (\mu - D)^{-1} p \\ v &= (\mu - D)^{-1} q. \end{aligned} \tag{1.3.7}$$

Using (1.3.7) for each eigenvalue λ_i of A , we find

$$Z^{-1} = \overbrace{\begin{bmatrix} -\text{diag}(p) \text{Cauchy}(D, \Lambda) \\ \mathbf{ones} \end{bmatrix}}^{\text{column eigenvectors}}, \quad \Pi Z = \overbrace{\begin{bmatrix} \text{Cauchy}(\Lambda, D) \text{diag}(q) & \mathbf{ones} \end{bmatrix}}^{\text{row eigenvectors}}. \tag{1.3.8}$$

The product of the matrices in (1.3.8) is not I , but it must be diagonal since the λ_j are simple eigenvalues.⁴

$$\begin{aligned} \Pi &= \Pi Z Z^{-1} = [\text{row eigenvectors}][\text{column eigenvectors}] \\ &= \begin{bmatrix} \text{Cauchy}(\Lambda, D) \text{diag}(q) & \mathbf{ones} \end{bmatrix} \begin{bmatrix} -\text{diag}(p) \text{Cauchy}(D, \Lambda) \\ \mathbf{ones} \end{bmatrix} \\ &= -\text{Cauchy}(\Lambda, D) \text{diag}(q) \text{diag}(p) \text{Cauchy}(D, \Lambda) + \mathbf{ones}. \end{aligned}$$

$$\text{By Theorem 1.3.4,} \quad q_i p_i = \prod_m (d_i - \lambda_m) \Big/ \prod_{k \neq i} (d_i - d_k). \quad \square$$

Now we are ready to specify the dual coordinates. Lemma 1.3.6 in our case gives

$$\begin{pmatrix} \Lambda_m & c_m \\ b_m^T & \delta_{m+1} \end{pmatrix} = \begin{pmatrix} -\text{diag}(c_m) \text{Cauchy}(\Lambda_m, \Lambda_{m+1}) \\ \mathbf{ones} \end{pmatrix} \Lambda_{m+1} \begin{pmatrix} -\text{diag}(c_m) \text{Cauchy}(\Lambda_m, \Lambda_{m+1}) \\ \mathbf{ones} \end{pmatrix}^{-1}.$$

Substituting this in (1.3.2) gives

$$x_{m+1} = g_{m+1} \Lambda_{m+1} g_{m+1}^{-1}$$

with

$$g_{m+1} = \begin{pmatrix} g_m & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} -\text{diag}(c_m) \text{Cauchy}(\Lambda_m, \Lambda_{m+1}) \\ \mathbf{ones} \end{pmatrix}.$$

Using Theorem 1.3.4, $\text{diag}(c_m) = -P_{m+1}(\Lambda_m) P'_m(\Lambda_m)^{-1} \text{diag}(b_m)^{-1}$, and so we get the g -recurrence

$$\begin{aligned} g_1 &= (1), \\ g_{m+1} &= \begin{bmatrix} -g_m \text{diag}(c_m) \text{Cauchy}(\Lambda_m, \Lambda_{m+1}) \\ \mathbf{ones} \end{bmatrix} \end{aligned} \tag{1.3.9}$$

$$= \begin{bmatrix} g_m P_{m+1}(\Lambda_m) P'_m(\Lambda_m)^{-1} \text{diag}(b_m)^{-1} \text{Cauchy}(\Lambda_m, \Lambda_{m+1}) \\ \mathbf{ones} \end{bmatrix}, \tag{1.3.10}$$

⁴If $A\tilde{u} = \lambda_i \tilde{u}$ and $\tilde{v}^T A = \lambda_j \tilde{v}^T$ for some $\tilde{u}, \tilde{v} \neq 0$ with $\lambda_i \neq \lambda_j$, then $\tilde{v}^T \tilde{u} = 0$.

and we find $x = g_n \Lambda_n g_n^{-1}$.

The recurrence (1.3.9) shows explicitly how Λ_m and Λ_{m+1} , i.e., $E(x_m)$ and $E(x_{m+1})$, determine the eigenvectors. Clearly (1.3.9) is simpler than (1.3.10), but we give preference to b_m over c_m to align our results with those of Kostant and Wallach.

We can put the parameters b_m together and define an invertible diagonal $\binom{n}{2} \times \binom{n}{2}$ matrix

$$b = \text{diag}(b_1, \dots, b_{n-1}) \in D\left(\binom{n}{2}\right),$$

where $D\left(\binom{n}{2}\right)$ is the group of invertible diagonal $\binom{n}{2} \times \binom{n}{2}$ matrices. The recurrence (1.3.10) constructs a matrix $g_n = g^{(b)} \in \text{GL}(n)$ given any $b \in D\left(\binom{n}{2}\right)$, and we have proved that

$$M_{\mathcal{R}}(n) \ni x \leftrightarrow \{ g_n \in \text{GL}(n) \mid g_n \Lambda_n g_n^{-1} \in M_{\mathcal{R}}(n) \text{ and } e_n^T g_n = \mathbf{ones} \} \leftrightarrow b$$

are bijections. We can use these bijections to define an action of $D\left(\binom{n}{2}\right)$ on $M_{\mathcal{R}}(n)$ by

$$\begin{aligned} D\left(\binom{n}{2}\right) \times M_{\mathcal{R}}(n) &\rightarrow M_{\mathcal{R}}(n) \\ b' \cdot x &= g^{(bb')} \Lambda_n (g^{(bb')})^{-1} \quad \text{if } x = g^{(b)} \Lambda_n (g^{(b)})^{-1}. \end{aligned}$$

This is a description of Kostant and Wallach's group action in [KW2, Theorem 5.9], which is revisited in Section 3 (see Remark 3.16) from a more natural point of view.

The conclusion (remember that we are still in the generic case) is that each choice of nonzero $s = (b_1^T, b_2^T, \dots, b_{n-1}^T)$ will determine a member of $M_{\mathcal{R}}(n)$, and different s 's yield different matrices in $M_{\mathcal{R}}(n)$. For each fixed s as $\mathcal{R}$ ranges over all generic $\mathcal{R}$'s we get a transverse slice of $M_{\Omega}(n)$. It seems a blemish that our dual coordinates b_m had to be non-zero; this will be removed naturally in the Lie format. The canonical coordinates will be angle coordinates q , while the non-zero coordinates will essentially appear as e^q . The non-zero coordinates do have the advantage of being single-valued on the fibres, though.

1.4 Complementary coordinates for the elementary conjugations.

For completeness's sake, we describe the dual coordinates that correspond to the elementary conjugations for generic $\mathcal{R}$.

Transposition. Let $b^T = (b_1^T, b_2^T, \dots, b_{n-1}^T)$ be the dual coordinates of $x \in M_{\mathcal{R}}(n) \subseteq M_{\Omega}(n)$. To find the dual coordinates of x^T , it is necessary to invoke the special diagonal matrices

$$\Sigma_m = -P_{m+1}(\Lambda_m) (P'_m(\Lambda_m))^{-1}, \quad 1 \leq m \leq n-1,$$

relating b_m to c_m which are given in Theorem 1.3.4, and also the diagonal matrices Π_m (appearing as Π in Lemma 1.3.6) which relate row and column eigenvectors. The diagonal matrices Σ_m and Π_m depend only on $\mathcal{R}$, not on b_m .

Lemma 1.4.1. Let the dual coordinates of x^T be $\tilde{b}^T = (\tilde{b}_1^T, \tilde{b}_2^T, \dots, \tilde{b}_{n-1}^T)$. Then, for $1 \leq m \leq n-1$,

$$\text{diag}(\tilde{b}_m) = \Pi_m \text{diag}(b_m)^{-1} \Sigma_m.$$

Proof. $x_m = g_m \Lambda_m g_m^{-1}$ implies

$$x_m^T = (g_m^{-1})^T \Lambda_m g_m^T = (g_m^{-1})^T \Pi_m \Lambda_m \Pi_m^{-1} g_m^T,$$

and the last row of $(g_m^{-1})^T \Pi_m$ is **ones** by (1.3.8). Transposition requires that the last row of $(\Pi_m^{-1} g_m^T \oplus 1) x_{m+1}^T ((g_m^{-1})^T \Pi_m \oplus 1)$ be equal to the last column of

$$(\Pi_m g_m^{-1} \oplus 1) x_{m+1} (g_m \Pi_m^{-1} \oplus 1) = \begin{pmatrix} \Lambda_m & \Pi_m c_m \\ b_m^T \Pi_m^{-1} & \delta_{m+1} \end{pmatrix},$$

therefore $\tilde{b}_m = \Pi_m c_m$, and the Lemma follows using $\text{diag}(b_m) \text{diag}(c_m) = \Sigma_m$. $\square$

Diagonal similarity. Let $\hat{x} = d x d^{-1}$. As usual, d_m denotes the leading principal submatrix of d , and so we denote the (m, m) th entry by $d(m)$. Let $\hat{b}^T = (\hat{b}_1^T, \hat{b}_2^T, \dots, \hat{b}_{n-1}^T)$ denote the dual coordinates of $d x d^{-1} = \hat{g}_n \Lambda_n \hat{g}_n^{-1}$. We note immediately that all g 's in (1.3.1) are normalized to have ones in the last row, so we cannot have $\hat{g}_m = d_m g_m$. To rectify this we define $\mathring{d}_m = d_m / d(m)$. In particular,

$$d_m x_m d_m^{-1} = \mathring{d}_m x_m \mathring{d}_m^{-1}.$$

Lemma 1.4.2. For generic $\mathcal{R}$,

$$\hat{b}_m = b_m d(m+1)/d(m), \quad 1 \leq m \leq n-1.$$

Proof. We have

$$\hat{x}_m = \mathring{d}_m x_m \mathring{d}_m^{-1} = \mathring{d}_m g_m \Lambda_m g_m^{-1} \mathring{d}_m^{-1},$$

so we calculate

$$\begin{aligned} (g_m^{-1} \mathring{d}_m^{-1} \oplus 1) \hat{x}_{m+1} (\mathring{d}_m g_m \oplus 1) &= (g_m^{-1} \mathring{d}_m^{-1} \oplus 1) d_{m+1} x_{m+1} d_{m+1}^{-1} (\mathring{d}_m g_m \oplus 1) = \\ &= d(m) \begin{pmatrix} g_m^{-1} \mathring{d}_m^{-1} & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \mathring{d}_m & 0 \\ 0 & \frac{d(m+1)}{d(m)} \end{pmatrix} x_{m+1} \left[d(m) \begin{pmatrix} g_m^{-1} \mathring{d}_m^{-1} & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \mathring{d}_m & 0 \\ 0 & \frac{d(m+1)}{d(m)} \end{pmatrix} \right]^{-1} = \\ &= \begin{pmatrix} d(m) I_m & 0 \\ 0 & d(m+1) \end{pmatrix} \begin{pmatrix} g_m^{-1} & 0 \\ 0 & 1 \end{pmatrix} x_{m+1} \left[\begin{pmatrix} d(m) I_m & 0 \\ 0 & d(m+1) \end{pmatrix} \begin{pmatrix} g_m^{-1} & 0 \\ 0 & 1 \end{pmatrix} \right]^{-1} = \\ &= \begin{pmatrix} d(m) I_m & 0 \\ 0 & d(m+1) \end{pmatrix} \begin{pmatrix} \Lambda_m & c_m \\ b_m^T & \delta_{m+1} \end{pmatrix} \begin{pmatrix} d(m)^{-1} I_m & 0 \\ 0 & d(m+1)^{-1} \end{pmatrix} = \\ &= \begin{pmatrix} \Lambda_m & \frac{d(m)}{d(m+1)} c_m \\ \frac{d(m+1)}{d(m)} b_m^T & \delta_{m+1} \end{pmatrix}, \end{aligned}$$

so that

$$\hat{c}_m = c_m (d(m)/d(m+1))$$

and

$$\hat{b}_m = b_m (d(m+1)/d(m)),$$

as claimed. $\square$

One may ask what exactly is preserved by elementary conjugations, and the answer is not Ritz values but rather all principal minor determinants. More precisely, a result due to Loewy [Lo, Theorem 1] is that under some non-degeneracy conditions two matrices have equal corresponding principal minors if and only if they are equivalent under an elementary conjugation.

There are $2^n - 1$ non-trivial principal minors, but only $n^2 - n + 1$ of them are independent, so the minors satisfy many relations, in contrast to what happens for Ritz values. While a full analysis of the problem is outside the scope of this article, we emphasize that this is again a prime example of the applicability of Lie- and representation-theoretic and geometric methods to a problem in matrix theory by exploiting its symmetry.

1.5 Genericity conditions

Consider the following problem: let $B_m \in M(m)$ be any matrix, and suppose we wish to find

$$B_{m+1} = \begin{pmatrix} B_m & c \\ b^T & \delta \end{pmatrix} \in M(m+1)$$

such that

$$\det(\lambda I_{m+1} - B_{m+1}) = \prod_{1 \leq i \leq m+1} (\lambda - \lambda_i) \quad (1.5.1)$$

for given $\lambda_1, \dots, \lambda_{m+1} \in \mathbb{C}$. (Recall that $\delta = \text{tr}(B_{m+1}) - \text{tr}(B_m)$ is fixed.) Equation (1.5.1) is an algebraic constraint on the $2m$ coordinates of b and c . The $2m$ coordinates must satisfy m polynomial equations, therefore under sufficiently general conditions we expect an m -dimensional set of solutions, while under degenerate conditions the dimension may increase, or there may be no solutions at all. Let us give several interpretations of this problem, and examine the role of each genericity condition. This will introduce useful notions from linear systems theory; this is another field not generally known to matrix theorists. These sections show why the strictly lower triangular part of x is not a viable choice of coordinates for x complementary to $\mathcal{R}$.

1.6 Observability and controllability

Consider the system of ordinary differential equations given by

$$\begin{aligned} \dot{x}(t) &= B_m x(t) + cu(t), \\ y(t) &= b^T x(t) + \delta u(t). \end{aligned} \quad (1.6.1)$$

This represents a continuous time-invariant linear system (SISO)⁵ with state $x(t) \in \mathbb{C}^m$, control $u(t) \in \mathbb{C}$, and output $y(t) \in \mathbb{C}$. Possibly abusing language,

The pair $\begin{pmatrix} B_m \\ b^T \end{pmatrix}$ is observable when $\text{rank}(b, B_m^T b, \dots) = m$.

The pair $\begin{pmatrix} B_m & c \end{pmatrix}$ is controllable when $\text{rank}(c, B_m c, \dots) = m$.

⁵SISO = Single Input, Single Output

The algebraic significance of observability and controllability will become clearer if we introduce the following terminology. Let $\mathbb{C}^m$ be the space of column vectors. A vector $v \in \mathbb{C}^m$ is called a cyclic vector for a matrix B_m if for any $w \in \mathbb{C}^m$ there exists a polynomial $f(x) \in \mathbb{C}[x]$ such that $f(B_m)v = w$. In this language, the system $\begin{pmatrix} B_m & c \end{pmatrix}$ is controllable if and only if c is a cyclic vector for B_m , and $\begin{pmatrix} B_m \\ b^T \end{pmatrix}$ is observable if and only if b is a cyclic vector for B_m^T . Matrix theorists would say that c is a cyclic vector for B_m if the minimal polynomial of c for B_m has (maximal) degree m : $f(B_m)c = 0$ for a nonzero polynomial f only if $\deg f \geq m$. The centralizer of an element x of a Lie algebra $\mathfrak{g}$ is the set of elements $\{y \in \mathfrak{g} \mid xy = yx\}$ that commute with x . (Lie algebras are briefly discussed elsewhere. Here one may read $\mathfrak{g} = M(n)$ and ignore the appellation. Those familiar with functional analysis or operator algebras will also recognize this as the definition of the commutant of $\{x\} \subseteq M(n) = \text{End}(\mathbb{C}^n)$.)

Theorem 1.6.2. A matrix B_m has a cyclic vector if and only if the centralizer of B_m coincides with the algebra $\{f(B_m) \mid f \in \mathbb{C}[x]\}$ of polynomials in B_m .

This property is known to Lie theorists as regularity (not to be confused with the property of being invertible—a matrix with only zero eigenvalues may well be regular in our sense). For matrix theorists, it is equivalent to being non-derogatory, i.e., the minimal polynomial equals the characteristic polynomial. Clearly the identity element is far from regular; a diagonal matrix is regular if and only if its diagonal entries are distinct.

Let $P_m(\lambda) = \det(\lambda I_m - B_m)$ and $P_{m+1}(\lambda) = \det(\lambda I_{m+1} - B_{m+1})$. Block elimination yields

$$P_{m+1}(\lambda) = P_m(\lambda)(\lambda - \delta - b^T(\lambda I_m - B_m)^{-1}c). \quad (1.6.3)$$

Assume (1.5.1), that is, that B_{m+1} has the specified eigenvalues.

Theorem 1.6.4. Given b , there exists a unique c such that (1.5.1) is satisfied if and only if $\begin{pmatrix} B_m \\ b^T \end{pmatrix}$ is observable. Given c , there exists a unique b such that (1.5.1) is satisfied if and only if $\begin{pmatrix} B_m & c \end{pmatrix}$ is controllable.

Proof. This is a variation on a standard problem in control theory. Re-write (1.6.3) as

$$b^T(\lambda I_m - B_m)^{-1}c = \lambda - \delta - \frac{P_{m+1}(\lambda)}{P_m(\lambda)}.$$

By construction (recall that δ is determined by $\delta = \text{tr}(B_{m+1}) - \text{tr}(B_m)$), the right-hand side is holomorphic at $\lambda = \infty$, and expanding both sides into power series gives

$$\sum_{k=1}^{\infty} \lambda^{-k} b^T B_m^{k-1} c = \sum_{k=1}^{\infty} \lambda^{-k} g_k$$

for some numbers $g_k \in \mathbb{C}$. Equating powers of λ , b and c must satisfy the (infinite) system of equations

$$b^T B_m^{k-1} c = g_k, \quad k = 1, 2, \dots$$

The condition that there exist a unique solution c is exactly that $(b^T \quad b^T B_m \quad \dots)^T$ have full column rank, i.e., observability.

The proof considering controllability is analogous. $\square$

Note. Condition (G2_m) holds exactly when $P_{m+1}(\lambda)$ and $P_m(\lambda)$ are relatively prime, in which case the right-hand side of

$$\sum_{k=1}^{\infty} \lambda^{-k} b^T B_m^{k-1} c = b^T (\lambda I_m - B_m)^{-1} c = \lambda - \delta - \frac{P_{m+1}(\lambda)}{P_m(\lambda)}.$$

is a rational function of degree m . In that case, if b and c constitute a solution, the Hankel matrix

$$\begin{pmatrix} b^T c & b^T B_m c & \dots \\ b^T B_m c & b^T B_m^2 c & \dots \\ \vdots & & \end{pmatrix} = \begin{pmatrix} b^T \\ b^T B_m \\ b^T B_m^2 \\ \vdots \end{pmatrix} (c \quad B_m c \quad \dots)$$

has rank m and so the system must be both observable and controllable.

Example. If B_m is an unreduced upper Hessenberg matrix, then the row $(0, \dots, 0, 1)$ always yields an observable system.

Example 1.6.5. Consider the case when B_m is a regular diagonal matrix (thus (G1_m) holds, and (G2_m) by assumption; we are not assuming (G1_{m+1})). Then (1.5.1) has solutions if and only if $\begin{pmatrix} B_m \\ b^T \end{pmatrix}$ is observable, if and only if $\begin{pmatrix} B_m & c \end{pmatrix}$ is controllable, if and only if all the entries of b and of c are non-zero. (See Theorem 1.3.4.)

If B_m is regular and semi-simple, but not diagonal, then it is still true that $\begin{pmatrix} B_m \\ b^T \end{pmatrix}$ is observable if and only if $\begin{pmatrix} B_m & c \end{pmatrix}$ is controllable: if $g_m^{-1} B_m g_m$ is diagonal, then this happens when $b^T g_m$, resp. $g_m^{-1} c$, has non-zero entries. (This is relevant to the matrix in (1.3.2).)

A matrix B_m has a cyclic vector (if and only if B_m^T has a cyclic vector), if and only if B_m is regular. So if B_m is not regular (in our sense), then the system $\begin{pmatrix} B_m \\ b^T \end{pmatrix}$ is never observable, nor is $\begin{pmatrix} B_m & c \end{pmatrix}$ ever controllable.

1.7 Beyond the generic case

The criterion in Example 1.6.5, that the entries of the row/column be non-zero, may be readily generalized to the case when B_m is regular, but (G1_m) fails to hold.

Lemma 1.7.1. Suppose that B_m is in the form

$$B_m = \text{diag}(J_{m_1}(d_1), \dots, J_{m_t}(d_t)),$$

where each Jordan block

$$J_{m_i}(d_i) = \begin{pmatrix} d_i & 0 & \dots \\ 1 & d_i & \dots \\ 0 & 1 & \dots \\ \vdots & & \end{pmatrix} \in M(m_i)$$

and $d_1, \dots, d_t$ are distinct. Then a row $(y_1, \dots, y_m)$ is observable if and only if each of the entries $y_{m_1}, y_{m_1+m_2}, \dots, y_m$ is nonzero, namely the last entry in each segment.

Sketch of proof. The matrix B_m^T is block-diagonal; let us write

$$B_m^T = J_{m_1}(d_1)^T \oplus \dots \oplus J_{m_t}(d_t)^T, \\ (y_1, \dots, y_m)^T = y^{(1)} \oplus \dots \oplus y^{(t)}, \quad y^{(i)} = (y_{k_i+1}, \dots, y_{k_i+m_i})^T, \quad k_i = \sum_{t' < i} m_{t'}$$

so that

$$B_m^T(y_1, \dots, y_m)^T = \bigoplus_{i=1}^m J_{m_i}(d_i)^T y^{(i)}.$$

We claim that $(y_1, \dots, y_m)^T$ is a cyclic vector for B_m^T if and only if $y^{(i)}$ is a cyclic vector for $J_{m_i}(d_i)^T$ for each $1 \leq i \leq t$. One way to see this is to recall that a vector y is cyclic for any matrix B_m if and only if $B'y = 0$ implies $B' = 0$, for every B' in the centralizer of B_m . Because our B_m^T is regular, its centralizer just consists of block-diagonal matrices $B' = J'_{m_1} \oplus \dots \oplus J'_{m_t}$, where each J'_{m_i} commutes with $J_{m_i}(d_i)^T$. This verifies our claim. This reduces the problem to calculating cyclic vectors for a Jordan block, which is straightforward. $\square$

Hence, if some generalized eigenvalue of a regular B_m has multiplicity, the set of rows making the system observable, and also the set of columns making the system controllable, is isomorphic to $(\mathbb{C}^\times)^t \times \mathbb{C}^{m-t}$. (And we recall that neither observability nor controllability is possible when B_m is not regular.)

We conclude with the observation that the geometric structure of the set of solutions for b and c satisfying (1.5.1), at least for regular B_m , depends only on how many Ritz values coincide, and not on the Ritz values themselves. This explains why transverse slices are possible when we restrict $\mathcal{R}$ to belong to a subset of Ritz values of fixed combinatorial type, such as the generic Ritz values introduced in Section 1.2.

2 Lie theory

For the reader who is not familiar with Lie theory and geometric terminology, we include an overview of the concepts necessary to present the results. Our purpose is not to give formal or abstract definitions, although we do so as necessary, but to paint a clear picture of the construction and how it relates to several important areas of mathematics. This section is independent of the rest of the paper, and may be ignored by the cognoscenti.

2.1 Basic geometry: integral curves on a vector field

This concept is constantly used in Section 3, every time a formula contains an expression like “ $\exp(q\xi)$.” It also paves the way for our discussion of integrable systems.

Start with a vector field ξ on M . (For concreteness, think of M as a smooth manifold, although everything goes through for complex or algebraic varieties, and, in particular, when we work with $M = M(n)$ our scalars will be in $\mathbb{C}$.) We have a tangent vector at each point of M . Starting at some $x \in M$, we may look for a curve that passes through x and is everywhere tangent to the vector field ξ :

$$\begin{aligned}\phi(0) &= x, \\ \frac{d}{dt}(\phi(t)) &= \xi(\phi(t)).\end{aligned}\tag{2.1.1}$$

By the theory of ordinary differential equations, there exists a unique solution for all t in some open interval containing 0, but there is, in general, no reason to expect a solution for all $t \in \mathbb{R}$.

Definition. If (2.1.1) does have a solution for all $x \in M$ and all $t \in \mathbb{R}$, one says, variously, that the flow defined by ξ is complete, or that ξ is complete, or that ξ is (globally) integrable.

For obvious reasons, the flow defined by (2.1.1) is called $\exp(t\xi)$, i.e., $\exp(t\xi)(x) = \phi(t)$, where $\phi(t)$ is defined as above. One can check that solutions to (2.1.1) satisfy $\exp((s+t)\xi) = \exp(s\xi)\exp(t\xi)$. The word “flow” is meant to suggest that as t changes all the points of M are smoothly displaced from their positions, like particles in a fluid.

Example. Since the vector field ξ has no singularities, intuitively there should be no obstruction to integrating it. If M is compact, then any vector field is complete (since there exists some $\varepsilon > 0$ such that a solution to (2.1.1) exists for $|t| < \varepsilon$ for any $x \in M$, and these patch together).

Example. As an example of what can happen when M is not compact, let $M = \mathbb{R}$ with coordinate x , and let $\xi = x^2 d/dx$. Then an integral curve of ξ through any point would satisfy $(d/dt)\phi(t) = \phi(t)^2$, whose solutions blow up. (The previous example shows that nothing bad happens as long as the integral curve remains bounded. If $\phi(0) \neq 0$, what happens is that we fall off the “end” of M in a finite amount of time. The same phenomenon may be seen with $\xi = d/dx$ and $M = (0, 1)$.)

Example. It may be argued that the last example is misleading: if we consider $\mathbb{R} \subseteq \mathbb{P}^1 = \mathbb{R} \cup \{\infty\}$, then

$$\frac{d}{dt}\phi(t) = \phi(t)^2 \rightsquigarrow \phi(t) = \frac{1}{\phi(0)^{-1} - t}$$

becomes a perfectly good flow, with a fixed point at $x = 0$. No such trick will enable one to integrate $x^3 d/dx$, though. The vector field $x^3 d/dx$ has a pole at $x = \infty$, and there is no way to embed $\mathbb{R}$ as a subset U of some manifold M and have a complete flow on M whose infinitesimal action on the subset U is $x^3 d/dx$.

2.2 Classical mechanics and Poisson geometry

The evolution of a mechanical system can be seen as the orbit of a point (initial state) under the action of time. In classical mechanics, the evolution of the system in time is determined by Hamilton's equations

$$\begin{aligned}\frac{dq_i}{dt} &= \frac{\partial H}{\partial p_i} \\ \frac{dp_i}{dt} &= -\frac{\partial H}{\partial q_i}.\end{aligned}\tag{2.2.1}$$

(for a particle in $\mathbb{R}^n$), where $q \in \mathbb{R}^n$ is position and $p \in \mathbb{R}^n$ is momentum, and the “Hamiltonian” function H is the energy of the particle.

We must keep track of time. Let $f = f(q, p)$ be a classical observable, which means any smooth function $\mathbb{R}^{2n} \rightarrow \mathbb{R}$. The function f does not depend on time, but let us write $f(t) = f(q(t), p(t)) : \mathbb{R}^{2n} \rightarrow \mathbb{R}$ for the observable resulting from picking a point, waiting for time t to elapse, and only then measuring the value of f . (In particular, $f(0) = f = f(q, p)$.)

Instead of considering just position or just momentum, we can re-write Hamilton's equations as

$$\frac{d}{dt}(f(t)) = \{H, f\}(t),\tag{2.2.2}$$

The right-hand side is the value at time t of the Poisson bracket

$$\{H, f\} = \sum_i \frac{\partial H}{\partial p_i} \frac{\partial f}{\partial q_i} - \frac{\partial H}{\partial q_i} \frac{\partial f}{\partial p_i}.\tag{2.2.3}$$

Physicists have concluded from staring at (2.1.1) and (2.2.2) that the Hamiltonian H generates time evolution: if the map $f \mapsto \{H, f\}$ is thought of as defining a vector field $\xi_H = \sum_i \frac{\partial H}{\partial p_i} \frac{\partial}{\partial q_i} - \frac{\partial H}{\partial q_i} \frac{\partial}{\partial p_i}$, then Hamilton's equations are satisfied if and only if the state of the system follows the flow of ξ_H :

$$f(t) = f \circ \exp(t\xi_H).\tag{2.2.4}$$

Therefore, time and time-evolution naturally appear as soon as one writes down the Hamiltonian. Any function H (thought of as “energy”) generates a complementary coordinate t (thought of as “time”), such that the system evolves in “time” so that energy is conserved. This is the essence of Hamiltonian geometry.

This formalism also goes through for any space M with a Poisson bracket, not just $\mathbb{R}^{2n}$. The appropriate generalization of (2.2.3) is

Definition. A manifold M is a Poisson manifold if the algebra $\mathcal{O}(M)$ of functions $M \rightarrow \mathbb{R}$ has a Poisson structure, i.e., there is a bracket

$$\{ , \} : \mathcal{O}(M) \otimes \mathcal{O}(M) \rightarrow \mathcal{O}(M)$$

making $\mathcal{O}(M)$ into a Lie algebra ($\{ , \}$ is bilinear, antisymmetric, and satisfies the Jacobi identity) and satisfying the Leibniz identity

$$\{f, gh\} = \{f, g\}h + g\{f, h\}.$$

Classical mechanics takes place on Poisson manifolds.

2.3 Integrable systems

Joseph Liouville concerned himself with characterizing those cases when explicit solutions to the equations (2.2.1) may actually be found. He proved this is the case when there are sufficiently many (independent) commuting Hamiltonians; this is known as complete integrability in the sense of Liouville. The idea is that for a completely integrable system we can (explicitly) find canonical coordinates.

We have seen in (2.2.4) that every function (“Hamiltonian”) on a phase space has a complementary coordinate associated with it, given by following some flow. Equation 2.2.2 says that a function f is constant along the flow of ξ_H if and only if $\{H, f\} = 0$. (In classical mechanics, such functions are called (first) integrals.) So, if $f_1, \dots, f_n$ are functions such that $\{H, f_i\} = 0$ and $\{f_i, f_j\} = 0$ for all i and j , then the trajectory of the system is contained in a level set of $(f_1, \dots, f_n)$. (Commutativity with respect to the Poisson bracket means that the flow corresponding to each function conserves all the other functions.)

Not every system has sufficiently many independent integrals of motion. (“Independent” means that their differentials are linearly independent (on a dense open subset of the phase space).) If there are enough independent first integrals which are simultaneously observable ($\equiv$ commutativity with respect to $\{ \ , \}$), then if the associated flows are complete we get a system of coordinates on the entire phase space given by the values of each of the functions together with the dual coordinates along the level sets (fibres!) given by following the flows.

Integrable systems are commonly defined in the case when M is a symplectic manifold, rather than in the more general case of a Poisson manifold. (A symplectic manifold is a manifold which has a closed non-degenerate 2-form; it is a very special kind of Poisson manifold.) Since we will later assert that Kostant and Wallach’s Gelfand–Zeitlin algebra defines an integrable system⁶ on $M(n)$, which is not symplectic, we give the more general definition.

First of all, what is the maximum possible number of independent commuting Hamiltonians? (For a symplectic M , this is $\frac{1}{2} \dim M$.) Let M be a Poisson manifold. The rank of the Poisson structure is defined to be the maximum possible number of linearly independent Hamiltonian vector fields at a point, i.e.,

$$\text{rank}\{ \ , \} = \max_{x \in M} \dim \langle (\xi_f)_x \mid f \in \mathcal{O}(M) \rangle.$$

One can show that any Poisson-commutative algebra of functions has dimension at most $\dim M - \frac{1}{2} \text{rank}\{ \ , \}$.

Definition 2.3.1. A Poisson manifold M of rank r together with a maximal Poisson-commutative algebra $A \subseteq \mathcal{O}(M)$ is a (completely) integrable Hamiltonian system if and

⁶This system, and its complete integrability, was already known to Thimm [T] and Guillemin–Sternberg [GS] in the 1980s.

only if

$$\dim A = \dim M - \frac{r}{2}.$$

Liouville showed [Li] how to construct canonical coordinates for an integrable system and solve Hamilton's equations. (A modern treatment may be found in [A]; for the generalization to the Poisson case, see, for example, [LMV].)

The coordinates dual to the functions are called angle coordinates.⁷ Denote them by φ_i ; we remark that once the angle coordinates are known, the system (2.2.1) is equivalent to

$$\begin{aligned}\frac{d}{dt}f_i(t) &= 0, \\ \frac{d}{dt}\varphi_i(t) &= \text{constant},\end{aligned}$$

which is trivial to integrate.

Note that the “canonical” coordinates do depend on the choice of Hamiltonians. Also, the complementary coordinates are measured from a basepoint, which must be specified (note that the level sets may not even be connected).

Example 2.3.2. An interesting system is a Lax pair

$$\frac{d}{dt}L(t) = [A, L],$$

where A and L are matrices. This differential equation describes an isospectral flow

$$L(t) = g(t)L(0)g(t)^{-1}$$

($g(t)$ and $A(t)$ are related via $dg/dt = Ag$), i.e., the eigenvalues of L are invariant over time. It follows that any functional f such that $f(gLg^{-1}) = f(L)$ is conserved; e.g., the coefficients of $\det(\lambda - L(t))$, or, alternatively, the functions $\text{tr}(L^m)$, are conserved quantities. Therefore, writing a system as a Lax pair exposes many integrals of motion (in fact, any completely integrable system can be written as a Lax pair, although constructing Lax pairs equivalent to integrable systems, and vice versa, is a far-from-trivial subject).

Not only is this an enormously successful method for actually solving various integrable systems (including various non-linear partial differential equations, such as KdV⁸), Lax pairs bring in Lie theory and geometry in a natural way. While no discussion of integrable systems can be complete without mentioning Lax pairs, the subject is too great to attempt a thorough treatment here; see [BBT] for an overview. For an application of Lax pairs to the topic of this paper, see [BP].

⁷Beware that the corresponding action coordinates are not usually the same as the particular functions used to specify an integrable system. In the case of generic Ritz values, the Ritz values themselves will be action coordinates (cf. [KW2, Theorem 5.23]), but this point is not needed in this paper and we will not pursue it.

⁸ $v_\tau = 6vv_z + v_{zzz}$

We adduce that Kostant and Wallach's theory should be thought of in this framework. There the phase space will be the space of all matrices, and the Hamiltonians—the conserved quantities—will be (certain symmetric functions of) the Ritz values; the fibres $M_{\mathcal{R}}(n)$ will be their level sets. Taking all of the Ritz values gives a maximal Poisson-commutative algebra of observables, and their number is exactly enough to make the system completely integrable.

3 Kostant–Wallach theory

The algebra of matrices $M(n)$ has a Poisson structure: let α_{ij} be the linear functional defined so that $\alpha_{ij}(x) = x_{ij}$, and let E_{ij} be the matrix with a 1 in the (i, j) th position and 0 elsewhere (so $\alpha_{ij}(E_{kl}) = \delta_{ij, kl}$ in terms of Kronecker's δ). Then

$$[E_{ij}, E_{kl}] = \delta_{jk}E_{il} - \delta_{il}E_{kj}$$

which specifies the Poisson structure as

$$\{\alpha_{ij}, \alpha_{kl}\} = \delta_{jk}\alpha_{il} - \delta_{il}\alpha_{kj}.$$

This extends naturally to define the Poisson bracket of any two polynomial, or even holomorphic, functions $M(n) \rightarrow \mathbb{C}$, because all such may be written in terms of the α_{ij} . The Leibniz rule yields

$$\{f, g\} = \sum_{ij, kl} \{\alpha_{ij}, \alpha_{kl}\} \frac{\partial f}{\partial \alpha_{ij}} \frac{\partial g}{\partial \alpha_{kl}}.$$

Keeping this Poisson structure in mind, motivation for Kostant and Wallach's theory may be found in the theory of completely integrable Hamiltonian systems.

Kostant and Wallach do not use any general theory in their original paper; the main actors there are the familiar algebra $M(n)$ of all $n \times n$ matrices, and the Lie group $\mathrm{GL}(n)$ of invertible $n \times n$ matrices, which acts on $M(n)$ via the adjoint representation

$$\mathrm{Ad}(g)x := gxg^{-1}. \quad (3.1)$$

Recall that we are interested in quantities conserved under the action of some group, and that we should look in advance for functions $M(n) \rightarrow \mathbb{C}$ that Poisson-commute. Even if we did not know about Ritz values, we might be led to consider them as follows.

If we were interested in studying $n \times n$ matrices up to similarity, we would be studying adjoint orbits, in other words, equivalence classes of matrices under similarity. A classical problem is to find numerical invariants of these adjoint orbits, namely all polynomial functions $f: M(n) \rightarrow \mathbb{C}$ (e.g., tr , det) such that $f(gxg^{-1}) = f(x)$ for all $x \in M(n)$ and $g \in \mathrm{GL}(n)$. The set of such functions may be denoted by

$$\mathrm{Pol}(M(n))^{\mathrm{GL}(n)} = \{ f \in \mathrm{Pol}(M(n)) \mid f \text{ is } \mathrm{GL}(n)\text{-invariant} \}.$$

The solution to this classical problem is that any such function is a symmetric polynomial in the roots of the characteristic polynomial, therefore is equal to a polynomial in $\mathrm{tr}(x^k)$,

$k = 1, \dots, n$.⁹ Since these functions are constant on adjoint orbits, not only do they Poisson-commute, but the corresponding vector fields are identically zero. However, using this observation, it follows by induction downwards on m that $\{\text{tr}(x_{m_1})^{k_1}, \text{tr}(x_{m_2})^{k_2}\} = 0$ for any m_1, k_1, m_2, k_2 (cf. [KW1, Proposition 2.1]).

Example. Let us write $\{\text{tr}(x_2), \text{tr}(x_3)^2\}$ out in coordinates. We can compute everything in terms of the linear functionals α_{ij} :

$$\{\text{tr}(x_2), \text{tr}(x_3)^2\} = \{\alpha_{11} + \alpha_{22}, \alpha_{11}^2 + \alpha_{22}^2 + \alpha_{33}^2 + 2\alpha_{12}\alpha_{21} + 2\alpha_{13}\alpha_{31} + 2\alpha_{23}\alpha_{32}\}.$$

If we use the Leibniz rule repeatedly and expand, we find that

$$\begin{aligned} \{\alpha_{11} + \alpha_{22}, \alpha_{11}^2 + 2\alpha_{12}\alpha_{21} + \dots\} &= \{\alpha_{11}, \alpha_{11}^2\} + 2\{\alpha_{11}, \alpha_{12}\alpha_{21}\} + \dots = \\ &= 2\alpha_{11}\{\alpha_{11}, \alpha_{11}\} + 2\{\alpha_{11}, \alpha_{12}\}\alpha_{21} + 2\alpha_{12}\{\alpha_{11}, \alpha_{21}\} + \dots = \\ &= 0 + 2\alpha_{12}\alpha_{21} - 2\alpha_{12}\alpha_{21} + \dots \end{aligned}$$

then we see that all terms cancel.

So all of the functions Poisson-commute, but the functions $\text{tr}(x_m)^k$ for $m < n$ are not Ad-invariant and their associated vector fields on $M(n)$ are non-zero. Someone looking for a Poisson-commutative algebra of functions on $M(n)$ might perhaps stumble upon Ritz values as a way to greatly enlarge $\text{Pol}(M(n))^{\text{GL}(n)}$.¹⁰

In any case, Kostant and Wallach begin by considering the algebra

$$J(n) = \text{Pol}(M(1))^{\text{GL}(1)} \text{Pol}(M(2))^{\text{GL}(2)} \dots \text{Pol}(M(n))^{\text{GL}(n)} \subseteq \text{Pol}(M(n)), \quad (3.2)$$

which is generated by the functions $\text{tr}((x_m)^k)$ for $m = 1, \dots, n$.¹¹ To make this notation clear, let us enumerate

$$f_1 = \text{tr}(x_1), \quad f_2 = \text{tr}(x_2), \quad f_3 = \text{tr}(x_2)^2, \quad \text{etc.} \quad (3.3)$$

Then a typical element of $J(n)$ looks like

$$\sum_{\alpha_i \geq 0} c_{\alpha_1, \alpha_2, \dots} f_1^{\alpha_1} f_2^{\alpha_2} \dots,$$

which maps $M(n) \rightarrow \mathbb{C}$. Since the f_i turn out to be algebraically independent, and they commute, $J(n)$ is isomorphic to a polynomial algebra $\mathbb{C}[f_1, \dots, f_{\binom{n+1}{2}}]$ in the variables f_i .

⁹As remarked in Example 2.3.2, any isospectral flow naturally conserves these quantities. To see that any Ad-invariant polynomial is a symmetric function of the roots of the characteristic polynomial, a quick way is to observe that any such polynomial is determined by its value on diagonal matrices.

¹⁰Kostant and Wallach explicitly mention in the abstract of [KW1] that they regard the algebra $J(n)$ generated by all the $\text{tr}(x_m)^k$ as a classical analogue of the Gelfand–Zeitlin algebra, which is a commutative (in the usual sense) subalgebra of the universal enveloping algebra of $M(n)$.

¹¹Taking linear combinations of products of functions on the submatrices may be new to many readers. A point of notation: we are implicitly using the truncation map $x \mapsto x_m$ to embed each $\text{Pol}(M(m))$

Proposition 3.4 ([KW1], Theorem 0.4). $J(n)$ is a (maximal) commutative subalgebra of $\text{Pol}(M(n))$. Furthermore, for any $f \in J(n)$, the Hamiltonian vector field ξ_f associated to f is globally integrable on $M(n)$, defining an action of $\mathbb{C}$ on $M(n)$.¹²

This richer structure (considering the set of all Ritz values of x simultaneously) now stands a chance of defining an integrable system on $M(n)$.

Proposition 3.5. Let $O_x \subseteq M(n)$ be an adjoint orbit¹³ consisting of regular¹⁴ elements. Then the Hamiltonians f_i , $1 \leq i \leq \binom{n}{2}$, form a completely integrable system on O_x . Moreover, the algebra $J(n)$ forms an integrable system on $M(n)$ in the sense of Definition 2.3.1.

This has a chance of being true because we have exhibited the right number of commuting Hamiltonians: $\dim O_x = n^2 - n$ for regular x , and the number of Hamiltonians is $1+2+\dots+(n-1) = n(n-1)/2$. Similarly, the Poisson rank of $M(n)$ is $n^2 - n$, and the total number of commuting Hamiltonians including also $f_{\binom{n}{2}+1}, \dots, f_{\binom{n+1}{2}} \in \text{Pol}(M(n))^{\text{GL}(n)}$ is $\binom{n+1}{2} = n^2 - (n^2 - n)/2$. (Things do not work out this nicely if, for instance, we replace $M(n)$ by the Lie algebra of symplectic matrices; see [C, Remark 1.7.2].)

Complete integrability requires that the Hamiltonians Poisson-commute and that they be independent. We have already mentioned the first condition; the question of independence leads to the notion of strong regularity. Kostant and Wallach give many equivalent characterizations of strong regularity. The ones relevant now are given by their Theorem 2.7 and the definition immediately preceding it:

Definition/Theorem. A matrix x is strongly regular if and only if the differentials $(df_i)_x$, $1 \leq i \leq \binom{n+1}{2}$, are linearly independent, if and only if the tangent vectors $(\xi_{f_i})_x$, $1 \leq i \leq \binom{n}{2}$, are linearly independent.

(The missing vector fields $\xi_{f_{\binom{n}{2}+1}}, \dots$, corresponding to the elements of $(\text{Pol } M(n))^{\text{GL}(n)}$, are zero, as explained before. Functions whose associated Hamiltonian vector fields are zero are called Casimir functions. One has to take them into account when dealing with integrable systems on a Poisson manifold, rather than the more familiar case of a symplectic manifold.)

In other words, a “strongly regular” matrix is a regular point of the function $x \mapsto (f_1(x), \dots, f_{\binom{n+1}{2}}(x))$. It does not mean what is sometimes called complete (or strong) regularity, that each x_m be invertible.

into $\text{Pol}(M(n))$, so each factor in (3.2) is a subalgebra of $\text{Pol}(M(n))$. This construction of a commutative algebra starting from a system of inclusions of subalgebras is associated with Gelfand–Zeitlin (a.k.a. Gelfand–Tsetlin), and is not meant to be intuitively obvious.

¹²Moreover, this action is given by a nice, explicit formula; see (3.7).

¹³We keep coming back to adjoint orbits. Their importance is that $M(n)$ is Poisson but not symplectic; the adjoint orbits are the symplectic leaves.

¹⁴An element $x \in M(n)$ is regular if and only if $\dim O_x = n^2 - n$. If x is not regular then $\dim O_x$ is strictly lower, and the Hamiltonians f_i have no chance of being independent there. Hence the hypothesis here that x be regular.

Kostant and Wallach prove [KW1, Theorem 2.3] that unit upper Hessenberg matrices are strongly regular, and, therefore, that the set of strongly regular matrices is a dense open subset of $M(n)$. This proves independence of the “Hamiltonians” f_i .

We are thus in the situation described before: any matrix will be described by its Ritz values, which specify a fibre $M_{\mathcal{R}}(n)$, and the complementary coordinates associated to the Ritz values (measured from a point on the fibre, which must be specified). The complementary coordinates will be angle coordinates for the integrable system.

The Gelfand–Zeitlin group A , central to their theory, is just the group obtained by integrating the vector fields corresponding to the functions $\text{tr}(x_m)^k \in J(n)$ for $1 \leq m \leq n-1$ and $1 \leq k \leq m$. This Lie group turns out not to be so mysterious; it is a commutative group, isomorphic to $\mathbb{C}^{\binom{n}{2}}$. If (3.3) are our chosen generators of $J(n)$, then a typical element of A can be written

$$a = \exp(q_1 \xi_{f_1}) \exp(q_2 \xi_{f_2}) \cdots \exp(q_{\binom{n}{2}} \xi_{f_{\binom{n}{2}}}) \quad (3.6)$$

where $q_i \in \mathbb{C}$. The reader unfamiliar with Lie groups may regard this expression as a rather formal way of keeping track of the coordinates q_i ; the group multiplication is given by

$$\left(\prod_i \exp(q_i \xi_{f_i}) \right) \left(\prod_i \exp(q'_i \xi_{f_i}) \right) = \prod_i \exp(q_i + q'_i) \xi_{f_i}.$$

The significance of the exponential map in Lie theory is that it relates maps on Lie groups with maps on Lie algebras. In this specific case, given any matrix $x \in M(n)$ and any $a \in A$, the matrix $a \cdot x \in M(n)$ is defined, and it is defined in terms of the vector fields ξ_{f_i} (which span a Lie algebra). This action is computed in [KW1], and we will use the results. In particular, we have the result, in compact form,

$$\exp \left(q \xi_{\text{tr}(x_m)^k} \right) \cdot x = \text{Ad} \left(\exp \left(-qk(x_m)^{k-1} \right) \oplus \mathbf{ones} \right) x, \quad (3.7)$$

giving an A -action on matrices.¹⁵ The key feature is that elements of A act by similarity transformations, and that those similarity transformations involve powers of leading principal submatrices of x . Moreover, $\mathcal{R}(a \cdot x) = \mathcal{R}(x)$ (“ A stabilizes the fibres $M_{\mathcal{R}}(n)$ ”).

When x is sufficiently generic, the fibre is a single A -orbit, that is, we have

$$M_{\mathcal{R}(x)}(n) = \{ a \cdot x \mid a \in A \}.$$

The $\binom{n}{2}$ parameters defining $a \in A$, together with the initial choice of x , induce a set of coordinates along the fibre, namely the q_j in (3.6).

In general, in view of (3.7), the orbit is given explicitly in terms of certain subgroups of $\text{GL}(n)$. To see which subgroups, note that the matrix $\exp((x_m)^{k-1})$ buried on the right-hand side of (3.7) is invertible and is a polynomial in x_m , and that successively

¹⁵The stated formula is an application of [KW1, Theorem 3.3]. Note the minus sign on the right side of (3.7).

applying (3.7) with various q and k , but keeping m fixed, results in $\text{Ad}(\exp(p(x_m)) \oplus \mathbf{ones})(x)$, where $p(x_m)$ can be any polynomial in x_m . Define

$$G_{x,m} = \{ g \in \text{GL}(m) \mid g \text{ is a polynomial in } x_m \}.$$

It will be convenient to consider $G_{x,m} \subseteq \text{GL}(m)$ as a subgroup of $\text{GL}(n)$ via the embedding $g \mapsto \text{diag}(g, \mathbf{ones})$.

Theorem 3.8 ([KW1], Theorem 3.7). The orbit $A \cdot x$ of (an arbitrary, not necessarily generic) matrix x is the image of the mapping

$$\begin{aligned} G_{x,1} \times G_{x,2} \times \cdots \times G_{x,n-1} &\rightarrow M(n) \\ (g(1), \dots, g(n-1)) &\mapsto \text{Ad}(g(1)) \cdots \text{Ad}(g(n-1))(x). \end{aligned} \quad (3.9)$$

This means that a general element in the A -orbit is obtained by performing a series of similarity transformations of a particular kind. Therefore, to describe $M_{\mathcal{A}}(n)$, we need to understand how it decomposes into A -orbits, and, to describe an A -orbit, we need to understand the kernel of (3.9).

These considerations lead to another characterization of strong regularity. The condition is that $\dim A \cdot x = \binom{n}{2}$ (the maximum possible).

Theorem 3.10. [[KW1], Theorem 3.14] Let x be strongly regular. Then the map

$$G_{x,1} \times \cdots \times G_{x,n-1} \rightarrow A \cdot x$$

is an algebraic isomorphism, so

$$A \cdot x \cong G_{x,1} \times \cdots \times G_{x,n-1},$$

where $G_{x,m}$ is the centralizer of x_m in $\text{GL}(m)$.

This reduces the description of the orbit of any strongly regular matrix to the description of the groups $G_{x,m}$, which can be done explicitly for any matrix. The only issue left is in picking a set of coordinates for $G_{x,m}$ that are somehow natural. (The group A was originally defined starting from particular symmetric functions of the Ritz values, but, for generic matrices, it will be more convenient to use instead the Ritz values themselves.)

Example 3.11. Let x_m be a regular semi-simple matrix, and suppose $x_m = g_m \Lambda_m g_m^{-1}$, where Λ_m is diagonal. Then its centralizer is $G_{x,m} = \{ g_m D g_m^{-1} \mid D \text{ is diagonal with non-zero entries} \} \cong (\mathbb{C}^\times)^m$. The parameters are the diagonal entries.

Example 3.12. Let x_m be any regular matrix, and suppose $x_m = g_m J_m g_m^{-1}$, where J_m is in Jordan canonical form. Then $G_{x,m} = \{ g_m D' g_m^{-1} \}$, where D' is a block-diagonal matrix whose blocks are invertible triangular Toeplitz matrices, one for each Jordan block. If the Jordan blocks are of sizes m_i with $m_1 + \cdots + m_t = m$, then $G_{x,m} \cong (\mathbb{C}^\times)^t \times \mathbb{C}^{m-t}$.

In integrable systems language, our Hamiltonians are independent, but $(f_1, \dots, f_{\binom{n+1}{2}})$ still has critical points—where the matrix is not strongly regular. Kostant and Wallach

give even more criteria for strong regularity, but we shall not need them. The point is that if x is not strongly regular, then describing $A \cdot x$ involves more than just calculating the groups $G_{x,m}$ and applying Theorem 3.10, even if each x_m happens to be regular.

Note. Even when x is strongly regular, if $(G2_m)$ is violated, meaning $E(x_m) \cap E(x_{m+1}) \neq 0$, then the description of $M_{\mathcal{R}(x)}(n)$ is complicated by the fact that A does not act transitively on the (strongly regular part $M_{\mathcal{R}(x)}^{\text{reg}}(n)$ of the) fibre, which breaks up into several (isomorphic) orbits. For example,

$$\begin{pmatrix} 0 & \dots & \dots \\ 1 & 0 & \dots \\ 0 & 1 & \dots \\ \vdots & & \end{pmatrix}$$

and its transpose belong to distinct orbits. A point in $M_{\mathcal{R}(x)}^{\text{reg}}(n)$ can be specified by indicating an element of A , the discrete data needed to specify a particular A -orbit, and a point in that orbit. Then, to define coordinates along the fibre, one needs to pick a representative of each orbit (note that there is a upper Hessenberg matrix in only one of the orbits); for example, the case $\mathcal{R} \equiv 0$ is worked out in [PS, C].

When x is generic, a more natural choice of functions would be the Ritz values $r_j(x)$ themselves. These are not globally defined functions, even restricted to the set of generic matrices (they are defined only on a covering). However, for any generic x , along with an ordering of each $E(x_m)$, it is possible to define vector fields η_j , $1 \leq j \leq \binom{n}{2}$, such that the action of $\mathbb{C}$ on $M_{\mathcal{R}(x)}(n)$ corresponding to the j th Ritz value is given by the action of $\exp(q\eta_j)$, $q \in \mathbb{C}$. What follows is the explicit expression of the associated similarities [see the proof of [KW2], Theorem 5.5]

Theorem 3.13.

$$\exp(q\eta_j) \cdot x = \text{Ad}(\gamma_j(e^{-q}))(x),$$

where

$$\gamma_j(e^{-q}) = \text{diag}(g_m \delta_l(e^{-q}) g_m^{-1}, \mathbf{ones}),$$

where, for $j = \binom{m}{2} + l$, $1 \leq l \leq m$, $g_m \in \text{GL}(m)$ is any matrix such that $x_m = g_m \Lambda_m g_m^{-1}$ and $\delta_l(e^{-q})$ is the $m \times m$ diagonal matrix

$$\text{diag}(\underbrace{1, \dots, 1}_l, e^{-q}, 1, \dots, 1).$$

Next we put together the similarities associated to all the eigenvalues of a submatrix x_m :

Corollary 3.14. Let $1 \leq m \leq n - 1$ and

$$a(m) = \prod_{\binom{m}{2} + 1 \leq j \leq \binom{m+1}{2}} \exp(q_j \eta_j), \quad q_j \in \mathbb{C}.$$

Then

$$a(m) \cdot x = \text{Ad} \left((g_m \text{diag}(e^{-q\binom{m}{2}+1}, \dots, e^{-q\binom{m+1}{2}}) g_m^{-1}) \oplus \mathbf{ones} \right) x, \quad (3.15)$$

where $g_m \in \text{GL}(m)$ is any matrix such that $x_m = g_m \Lambda_m g_m^{-1}$.

Proof. Apply Theorem 3.13, noting that the same g_m works for each $j = \binom{m}{2}+1, \dots, \binom{m+1}{2}$. $\square$

Remark 3.16. A generic fibre $M_{\mathcal{R}}(n)$ is a single orbit, and the element $\exp(q_1 \eta_1) \cdots \exp(q_{\binom{n}{2}} \eta_{\binom{n}{2}})$ of A acts as the identity on the fibre if and only if each $q_j \in 2\pi i \mathbb{Z}$. (See [KW2], Theorem 5.9.) Corollary 3.14 shows that the entries of the diagonal matrices D in Example 3.11 are, in fact, the coordinates e^{-q_j} dual to the Ritz values. The condition that the coordinates b_m^T and c_m not vanish is filled automatically here by the exponentials. Geometrically, a generic fibre is an $\binom{n}{2}$ -dimensional torus, because it is isomorphic to a product of $\binom{n}{2}$ copies of the multiplicative group $\mathbb{C}^\times$.

Remark. The coordinates introduced in Example 3.12 are a direct generalization, but it would be interesting to check whether they satisfy some nice properties analogous to the generic case.¹⁶

Finally, Kostant and Wallach define the coordinates s_j on a generic $M_{\mathcal{R}}(n)$ by picking an initial point. Recall (Theorem 1.1.3) that $M_{\mathcal{R}}(n)$ contains a unique unit upper Hessenberg matrix y . Then s_j is defined by

$$s_j \left(\exp(q_1 \eta_1) \cdots \exp(q_{\binom{n}{2}} \eta_{\binom{n}{2}}) \cdot y \right) = e^{-q_j}$$

(so $s_j(y) = \mathbf{ones}$).

3.1 Relation to arrow coordinates

We now prove Claim 1.3.3, that (s_j) above are identical to the arrow coordinates $(b_1^T, \dots, b_{n-1}^T)$ defined in Section 1.3, at the end of the matrix development.

Claim. If

$$x = \tilde{a}(1) \cdots \tilde{a}(n-1) \cdot y$$

with

$$\tilde{a}(m) = \prod_{\binom{m}{2}+1 \leq j \leq \binom{m+1}{2}} \exp(-q_j \eta_j),$$

then (1.3.2) holds with $b_m^T = (e^{q\binom{m}{2}+1}, \dots, e^{q\binom{m+1}{2}})$.

¹⁶In the generic case the “nice property” is that the diagonal entries in D in Example 3.11 are exponentials of angle coordinates. This says something about the symplectic geometry of generic matrices. Generalizing this invokes the geometry of certain less generic strata of the space of strongly regular ma-

Proof. Let $g_m \in \text{GL}(m)$, $1 \leq m \leq n-1$, be the special matrices defined by applying the procedure described in Section 1.1 to the unit Hessenberg matrix y . We shall need the fact

$$y_m = g_m \Lambda_m g_m^{-1} \quad (3.1.1)$$

as well as the normalization

$$e_m^T g_m = \mathbf{ones}, \quad e_m^T = (0, \dots, 0, 1), \quad (3.1.2)$$

i.e., the last row of g_m is **ones**.

By Corollary 3.14, $x = \text{Ad}(g(1)) \cdots \text{Ad}(g(n-1))y$ with

$$g(m) = g_m \text{diag}(b_m)^{-1} g_m^{-1} \oplus \mathbf{ones} \in G_{y,m}. \quad (3.1.3)$$

Observe that conjugation by an element of $G_{y,m}$ leaves y_m fixed, so

$$\begin{aligned} (\text{Ad}(g(n-1))y)_{n-1} &= y_{n-1}, \\ (\text{Ad}(g(n-2)) \text{Ad}(g(n-1))y)_{n-2} &= \text{Ad}(g(n-2)_{n-2})y_{n-2} = y_{n-2}, \end{aligned}$$

etc., and we have

$$x_m = \text{Ad}(g(1)_m \cdots g(m-1)_m)y_m, \quad (3.1.4)$$

$$x_{m+1} = \text{Ad}(g(1)_{m+1} \cdots g(m)_{m+1})y_{m+1}. \quad (3.1.5)$$

But (3.1.1) and (3.1.4) imply that $x_m = Z_m \Lambda_m Z_m^{-1}$ with

$$Z_m = g(1)_m \cdots g(m-1)_m g_m, \quad (3.1.6)$$

whence

$$\begin{aligned} (Z_m^{-1} \oplus 1)x_{m+1}(Z_m \oplus 1) &= \text{Ad}((Z_m^{-1} \oplus 1)g(1)_{m+1} \cdots g(m)_{m+1})y_{m+1} \\ &= \text{Ad}((g_m^{-1} \oplus 1)g(m)_{m+1})y_{m+1}, \quad \text{by substituting (3.1.6),} \\ &= \text{Ad}((g_m^{-1} \oplus 1)(g_m \oplus 1)(\text{diag}(b_m)^{-1} \oplus 1)(g_m^{-1} \oplus 1))y_{m+1}, \quad \text{by (3.1.3),} \\ &= (\text{diag}(b_m)^{-1} \oplus 1)(g_m^{-1} \oplus 1)y_{m+1}(g_m \oplus 1)(\text{diag}(b_m) \oplus 1). \end{aligned} \quad (3.1.7)$$

Due to our normalization of g_m , we have (suppressing irrelevant entries)

$$\begin{pmatrix} g_m^{-1} & 0 \\ 0 & 1 \end{pmatrix} y_{m+1} \begin{pmatrix} g_m & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} g_m^{-1} & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} y_m & * \\ e_m^T & * \end{pmatrix} \begin{pmatrix} g_m & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} g_m^{-1} y_m g_m & * \\ e_m^T g_m & * \end{pmatrix} = \begin{pmatrix} \Lambda_m & * \\ \mathbf{ones} & * \end{pmatrix},$$

therefore (3.1.7) becomes

$$\begin{pmatrix} \text{diag}(b_m)^{-1} & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \Lambda_m & * \\ \mathbf{ones} & * \end{pmatrix} \begin{pmatrix} \text{diag}(b_m) & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} \Lambda_m & * \\ b_m^T & * \end{pmatrix}.$$

□

trices, where the eigenvalues are allowed to coalesce.

3.2 Coordinates along a non-generic fibre

As an illustration that the ideas of Section 3 can be generalized to describe any fibre, we consider a case studied by Mark Colarusso [C]. He suggests looking at the set of matrices satisfying

- x_m is regular, $1 \leq m \leq n$.
- $(G2_m)$, $1 \leq m \leq n - 1$.

Any such matrix is strongly regular, and, moreover, the second condition further implies that $M_{\mathcal{R}(x)}^{\text{reg}}(n)$ is a single A -orbit. (This is the largest set of matrices that can be specified by naming a fibre and an element of A .)

The disadvantage of relaxing $(G1_m)$ is that there will no longer be a global set of complementary coordinates, because the geometry of the fibre will vary when the multiplicity of an eigenvalue changes. A proper generalization of Kostant and Wallach's results in [KW2] would consider the geometry of the space of strongly regular matrices (satisfying $(G2_m)$, for simplicity) such that the generalized eigenvalues of x_m have given multiplicity. However, since we avoided the technical complication of constructing global coordinates by considering the fibres individually, we can allow ourselves to examine a single fibre in this slightly less generic case.

The answer is given by Example 3.12, and the corresponding arrow coordinates may be computed as follows. Recall that any eigenvector of x_m has non-zero last entry.

Claim 3.2.1. If the Jordan form of x_m is $J_{m_1}(\mu_1) \oplus \cdots \oplus J_{m_t}(\mu_t)$, where each $J_{m_l}(\mu_l)$ is a lower Jordan block, then there exists $g_m \in \text{GL}(m)$ such that $x_m = g_m J_m g_m^{-1}$ and the last row of g_m is

$$(\underbrace{0, 0, 0, \dots, 1}_{m_1}, \underbrace{0, \dots, 1}_{m_2}, \dots). \quad (3.2.2)$$

Given m , let g_m be as in Claim 3.2.1. Then the arrow coordinates b_m^T are given by the first m entries of the bottom row of $\text{diag}(g_m^{-1}, 1)x_{m+1}\text{diag}(g_m, 1)$. The coordinates of a unit upper Hessenberg matrix are (3.2.2), and this coincides with the previous construction in case x is a generic matrix.

References

- [A] V. I. Arnol'd. Mathematical Methods of Classical Mechanics, volume 60 of Graduate Texts in Mathematics. Springer-Verlag, New York, 1997. Translated from the 1974 Russian original by K. Vogtmann and A. Weinstein, Corrected reprint of the second (1989) edition.
- [BBT] Olivier Babelon, Denis Bernard, and Michel Talon. Introduction to Classical Integrable Systems. Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge, 2003.

- [BP] Roger Bielawski and Victor Pidstrygach. Gelfand–Zeitlin actions and rational maps. arXiv:math.SG/0612365.
- [C] Mark Colarusso. The Gelfand–Zeitlin Algebra and Polarizations of Regular Adjoint Orbits for Classical Groups. PhD thesis, University of California, San Diego, 2007.
- [GS] V. Guillemin and S. Sternberg. The Gelfand–Zeitlin system and quantization of the complex flag manifolds. *J. Funct. Anal.*, 52(1):106–128, 1983.
- [KW1] Bertram Kostant and Nolan Wallach. Gelfand–Zeitlin theory from the perspective of classical mechanics. I. In *Studies in Lie theory*, volume 243 of *Progr. Math.*, pages 319–364. Birkhäuser Boston, Boston, MA, 2006.
- [KW2] Bertram Kostant and Nolan Wallach. Gelfand–Zeitlin theory from the perspective of classical mechanics. II. In *The unity of mathematics*, volume 244 of *Progr. Math.*, pages 387–420. Birkhäuser Boston, Boston, MA, 2006.
- [Li] J. Liouville. Sur l’intégration des équations différentielles de la Dynamique. *Journal de Mathématiques Pures et Appliquées*, XX, Mai 1855.
- [LMV] Camille Laurent, Eva Miranda, and Pol Vanhaecke. Action-angle coordinates for integrable systems on Poisson manifolds. arXiv:0805.1679 [math.SG].
- [Lo] Raphael Loewy. Principal minors and diagonal similarity of matrices. *Linear Algebra Appl.*, 78:23–64, 1986.
- [P] Beresford N. Parlett. *The Symmetric Eigenvalue Problem*, volume 20 of *Classics in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1998. Corrected reprint of the 1980 original.
- [PS] Beresford Parlett and Gilbert Strang. Matrices with prescribed Ritz values. *Linear Algebra Appl.*, 428(7):1725–1739, 2008.
- [T] A. Thimm. Integrable geodesic flows on homogeneous spaces. *Ergodic Theory and Dynamical Systems*, 1:495–517, 1980.